

Ophiura sarsii sarsii (Echinodermata, Ophiuroidea) from the Late Pliocene Hachioji Formation in Niigata Prefecture, Central Japan

YOSHIAKI ISHIDA and YOSHITAKA KURITA

Hitotsubashi Senior High School, 1-12-13 Higashikanda, Chiyoda-ku, Tokyo 101-0031, Japan

Shiozawa Primary School, 1550-1, Oaza Shiozawa, Shiozawa-machi, Minami Uonuma-gun, Niigata 949-6408, Japan

Received 11 February 1998; Revised manuscript accepted 20 May 1998

Abstract. Specimens of the Recent species *Ophiura sarsii sarsii* Lütken, collected from the Late Pliocene Hachioji Formation in Kashiwazaki City, Niigata Prefecture, central Japan, represent the first record of this species from the Pliocene, although this species has been found from the Middle-Late Miocene. The fossil specimens have been compared and contrasted with related species using morphometric characteristics of the disk, basal arm portions and radial shields.

Key words: Hachioji Formation, Late Pliocene, *Ophiura sarsii sarsii*, Ophiuroidea

Introduction

Ophiura sarsii sarsii is an extant, circumpolar species in the Northern Hemisphere, occurring in high-density populations in the upper bathyal zone surrounding northern Japan (Fujita and Ohta, 1989). Fossil ophiuroids assigned to *Ophiura s. sarsii* have been found in the Pleistocene from off northern Norway (Jensen and Thomsen, 1987), from southern Norway (Bjørlykke, 1898), from the Middle Pleistocene Ichijuku Formation (Kazusa Group) in Chiba Prefecture (Ishida and Inoue, 1993, 1995), from the Plio-Pleistocene Hijikata Formation in Shizuoka Prefecture (Ishida et al., 1996), from the Late Miocene Ogawa Formation in Nagano Prefecture (Ishida, Kurita et al., 1997; Ishida et al., 1998), from the Late Miocene Hongo Formation in Yamagata Prefecture (Ishida, Tokairin et al., 1997), and from the Middle-Late Miocene Wakkanai Formation in Hokkaido (Ishida and Fujita, 1998).

We have recently discovered fossil ophiuroids in the Late Pliocene Hachioji Formation, Kashiwazaki City (Ishida, Kurita et al., 1997), which proved to be assignable to *Ophiura s. sarsii*, based on a detailed morphological analysis. This paper describes *Ophiura s. sarsii* from the Pliocene in detail and discusses the comparison of the fossils with related species, which are important for its identification.

Locality and age

Thirty-five individuals were collected from a cliff outcrop of the Hachikoku Oil Field, about two kilometers southeast of Nagatori Station on the Japan Railways Shin-etsu Line in the district of Kashiwazaki City, Niigata Prefecture in May 1995

(Figure 1). The Pliocene and Pleistocene in this area comprise, in ascending order, the Hododaira Formation, the Hachikokusan Formation, the Suganuma Formation, the Hachioji Formation and the Uonuma Formation (Yasui et al., 1983; Kobayashi et al., 1989). The strata in this cliff are




Figure 1. Sampling locality in the Hachikoku Oil Field. Part of the "Kashiwazaki" 1:50,000 topographic map by the Geographical Survey Institute. ×: Sampling locality.

composed of massive sandy siltstone assigned to the middle part of the Hachioji Formation (Yasui *et al.*, 1983; Kobayashi *et al.*, 1989).

The age of the Hachioji Formation is inferred to be Late Pliocene from the following studies. The formation was correlated with the Pliocene Nishiyama Formation of the standard succession in the Niigata area on the basis of tephra-stratigraphy and foraminiferal biostratigraphical data (Kobayashi *et al.*, 1989). From an analysis of the stratigraphic sequence of the Niigata sedimentary basin, the Hachioji Formation falls into the Late Pliocene (Arato, 1997). The Hachioji Formation was correlated lithologically with the Asojima Formation in the Yoneyama area (Kobayashi *et al.*, 1989). Fission track ages of 3.24 and 2.91 Ma have been obtained for the Asojima Formation (Unpublished data of Muramatsu, in Kobayashi *et al.*, 1989).

Fossil specimens of *Ophiura s. sarsii* have been described from the Middle Pleistocene, the Plio-Pleistocene boundary and the Late Miocene (Ishida and Inoue, 1993; Ishida *et al.*,

1996, 1998), and also from the Middle-Late Miocene (Ishida and Fujita, 1998), but this is the first report of the species from the Pliocene.

Systematic description

Family Ophiuridae Lyman, 1865
Subfamily Ophiurinae Lyman, 1865
Genus *Ophiura* Lamarck, 1816

Ophiura sarsii sarsii Lütken, 1855

Figures 2, 3

Ophiura sarsii Lütken, 1855, p. 101; Clark, 1911, p. 37; Matsumoto, 1917, p. 272, fig. 74; Mortensen, 1927, p. 238, figs. 128-1, 2; Berry, 1934, p. 98, pls. 5, 6; D'yakonov, 1954, p. 98, fig. 35; Irimura, 1990, p. 98; Ishida and Inoue, 1993, p. 104, pls. 1-3; Ishida *et al.*, 1996, p. 67-69, fig. 3; Ishida *et al.*,




Figure 2. Camera lucida drawings of the figured specimens of *Ophiura sarsii sarsii* Lütken, 1855 from the Hachioji Formation. **A.** Dorsal view of disk and proximal arm; **B.** Ventral view, proximal to mid arm; **C.** Dorsal view, proximal to mid arm; **D.** Latero-ventral view, proximal to mid arm; **E.** Partial ventral view of disk; **F.** Lateral view of lateral arm plate (inside); **G.** Entire animal, showing ventral side (vertebral ossicles showing dorsal side). Abbreviation: 1, Lateral arm plate (ventral view); 2, Arm spine; 3, Ventral arm plate; 4, Lateral arm plate (lateral side); 5, Vertebral ossicle (dorsal side); 6, A part of genital plate; 7, Oral plate; 8, Adoral plate; 9, Oral shield; 10, Disk scale; 11, Primary scale; 12, Radial shield; 13, First dorsal arm plate; 14, Dorsal arm plate; 15, First ventral arm plate; 16, Arm comb plate; 17, Socket of arm spine. Scale bars equal 1 mm.




Figure 3. Fossil *Ophiura sarsii sarsii* Lütken, 1855 from the Hachioji Formation. **A.** Dense occurrence of fossils; **B.** Small individuals; **C.** Dorsal view of a complete specimen; **D.** Free arm, proximal part showing inside of ventral arm plates and lateral arm plates, distal part showing dorsal side of vertebral ossicles; **E.** Free arm, latero-ventral view; **F.** Ventral view of disk and proximal arms; **G.** Dorsal view of disk and proximal arm. Morphological explanation: 1, Inside of ventral arm plate; 2, Inner ventral side of lateral arm plate; 3, Dorsal view of vertebral ossicle; 4, Lateral side of lateral arm plate; 5, Arm spine; 6, Ventral view of vertebral ossicle; 7, A part of radial shield; 8, Central plate; 9, Scale; 10, Lateral side of lateral arm plate; 11, Dorsal arm plate; 12, Disk margin.

1998, p. 10-12, figs. 3, 4.

Materials.—Thirty-five specimens from the Hachioji Formation, three of which are illustrated here, are housed at the Municipal Nagaoka Science Museum, Niigata Prefecture

(Gf8-10).

Measurements.—Measurements are based on 31 specimens; disk diameter ranges from 6.0-14.1 mm; mean, 9.7 mm; median, 9.9 mm.

Description.—Disk circular in outline, low and flat, covered with small, flat and imbricated scales. Primary scales fairly large and circular. Radial shields oval, about twice as long as wide, separated from each other and about half as long as disk radius. Comb plates elliptical. Oral shields about one third of disk radius, pentagonal with rounded distal borders, with a pointed corner proximally and slightly longer than wide. Adoral plates slender, rectangular, in contact with each other at adoral margin. Oral plates fairly long, rectangular with a pointed corner proximally, in contact with each other at adoral side. Genital plates slender and long. First 4 or 5 arm segments insert laterally into disk. Arms

bent gradually on bedding plane, more than three times as long as disk diameter. Arms flattened, much wider than high, rather wide at base, tapering gradually. Dorsal arm plates well developed, rectangular, wider than long, with median keel, successive plates broadly in contact. First dorsal arm plates triangular. Ventral arm plates triangular, about 2–3 times as wide as long. First ventral arm plates trapezoidal. Lateral arm plates well developed, separated by dorsal arm plates, but in contact ventrally. Arm spines long and tapering, about twice as long as arm segment proximally and almost equal to length of arm segment at mid-arm, three in number, adpressed or often somewhat




Figure 4. Size frequency distribution of Recent *Ophiura sarsii sarsii*, *Ophiura leptoctenia*, *Ophiura kinbergi* (A) and fossil *Ophiura sarsii sarsii* from the Hachioji Formation (B). Data for these Recent specimens are from the specimens stored at the National Science Museum, Tokyo (*Ophiura s. sarsii*, NSMT-E 1608, Wakasa Bay, Fukui Prefecture, 270 m depth; 1414, Toyama Bay, Toyama Prefecture, ca. 200 m depth; *Ophiura leptoctenia*, NSMT-E 1987, Off Otsuchi, Iwate Prefecture, 1,038–1,055 m depth; *Ophiura kinbergi*, NSMT-E 0670, Amakusa Isls., Kumamoto Prefecture, ca. 30 m depth).




Figure 5. The relationship between disk diameter and basal arm width of Recent *Ophiura sarsii sarsii*, *Ophiura leptoctenia* and fossil *Ophiura sarsii sarsii* from the Hachioji Formation. Solid lines show linear regression and broken lines show 95% confidence limits. Each regression line is statistically significant ($r=0.99$, $p<0.05$ for living *O. s. sarsii* (Oss), $r=0.95$, $p<0.05$ for *O. leptoctenia* (Ol) and $r=0.97$, $p<0.05$ for fossil specimens of *O. s. sarsii* from the Hachioji Formation (H)). There is a statistically significant difference on intercept ($p<0.05$) and regression coefficient ($p<0.05$) between the two regression lines (Oss, Ol). The fossil specimens safely fall into the confidence interval of Recent *O. s. sarsii*, but not into that of *O. leptoctenia*. Data for Recent *O. s. sarsii* and *O. leptoctenia* are from specimens stored at the National Science Museum, Tokyo (*O. s. sarsii* NSMT-E 0568, Wakasa Bay, 275 m depth; 1414, 1608, 1609, Wakasa Bay, 240 m depth; *O. leptoctenia* NSMT-E 1987).

detached. Vertebral ossicles, triangular dorsally, with a pointed distal corner.

Remarks.—Morphologically the Hachioji specimens have much in common with Recent *O. s. sarsii*, which is related to *Ophiura s. vadicola*, *Ophiura kinbergi* and *Ophiura leptoctenia*. These four (sub)species can be distinguished mainly on the basis of the shape of arm comb papillae. Although the arm comb papillae were not discernible in the Hachioji fossils, they have been identified as *Ophiura s. sarsii* for the following reasons. Recent *O. s. sarsii* is larger than *O. kinbergi*, while the Hachioji specimens are similar to *O. s. sarsii* in size (Figure 4). Recent *O. s. sarsii* has a larger ratio of basal arm

width to disk diameter than has *O. leptoctenia*, while the Hachioji specimens are close to *O. s. sarsii* in this ratio: the values are 0.24 for *O. s. sarsii*, 0.12 for *O. leptoctenia* and 0.22 for the Hachioji specimens (Figure 5). Recent specimens of *O. s. sarsii* possess a smaller width to length ratio for radial shields than *O. s. vadicola* (the ratio is 0.75 for *O. s. vadicola* and 0.54 for *O. s. sarsii*), while the Hachioji specimens are similar to *O. s. sarsii* (the ratio is 0.58) (Figure 6). Recent *O. s. sarsii* has longer radial shields than those of Recent *O. s. vadicola* (the length is 0.5 times disk radius in *O. s. sarsii*, and 0.4 times disk radius in *O. s. vadicola* based on the same samples as Figure 6), while the Hachioji specimens are similar to *O. s. sarsii*. The body size of the Hachioji specimens is similar to that of specimens from the Lower Pleistocene Ichijuku Formation (mean disk diameter 9.6 mm) (Ishida and Inoue, 1993), but is larger than that of specimens from the Plio-Pleistocene Hijikata Formation (mean disk diameter 8.2 mm) (Ishida *et al.*, 1996).




Figure 6. The relationship between width and length of radial shields in Recent *Ophiura sarsii vadicola*, *Ophiura sarsii sarsii* and a fossil specimen from the Hachioji Formation. The value at each point for *O. s. vadicola* and *O. s. sarsii* indicates the mean value measured from five radial shields in the specimen. The value for the fossil specimen indicates the mean value measured from three radial shields in the specimen. Solid lines show linear regression and broken lines show 95% confidence limits. Each regression line is statistically significant ($r=0.97$, $p<0.05$ for living *O. s. vadicola*, $r=0.95$, $p<0.05$ for *O. s. sarsii*). There is a statistically significant difference on intercept ($p<0.05$) and regression coefficient ($p<0.05$) between the two regression lines. The fossil specimen safely falls into the confidence interval of Recent *O. s. sarsii*, but not into that of *O. s. vadicola*. Data for Recent specimens of *O. s. vadicola* and *O. s. sarsii* are from the specimens stored at the National Science Museum, Tokyo (*O. s. vadicola* NSMT-E1821, *O. s. sarsii* NSMT-E1609, 0568).

Mode of occurrence and paleoenvironment

Dense aggregations of fossilized ophiuroids were found in sandy siltstone layers. Many of their arms and disks are still attached, and most individuals (85%) lie dorsal side up on the bedding plane (Figure 7). This suggests the assemblage is




Figure 7. Sketch illustrating the dense occurrence of fossil *Ophiura sarsii sarsii* in sandy siltstone matrix from the Hachioji Formation.

Table 1. List of molluscan fossils associated with fossil ophiuroids. N; Numbers. For Recent species, depth and latitudinal range are given, according to Habe (1977), Higo (1973) and Kuroda and Habe (1952).

Species	N	Depth (m)	Latitude (°N)
Pelecypoda			
<i>Saccela confusa</i> (Hanley)	4	10-60	-35
<i>Nuculana (Thestylea) yokoyamai</i> (Kuroda)	3	50-450	32-43
<i>Acila</i> sp.	1		
<i>Yoldia (Cnesterium) notabilis</i> Yokoyama	4	50-100	32-48
<i>Portlandia (Megayoldia) thraciaeformis</i> (Storer)	5	25-550	37-70
<i>Limopsis (Empleconia) cumingi</i> A. Adams	4	30-300	31-36
<i>Anadara (Scapharca) ommaensis</i> Otuka	3		
<i>Clementia vatheleti</i> Mebille	5	0-140	33-41
<i>Solamen spectabilis</i> (A. Adams)	3	30-300	30?-42
<i>Nemocardium (Keenaea) cf. samarangae</i> (Makiyama)	3	50-300	30-42
<i>Felaniella usta</i> (Gould)	5	10-150	33-45
<i>Megacardia ferruginosa</i> (A. Adams and Reeve)	4	10-120	31-42
<i>Macoma calcarea</i> (Gmelin)	2	0-1000	33-72
<i>Thracia cf. kakumana</i> Yokoyama	1		
Gastropoda			
<i>Turritella (Neohaustator) saishuensis</i> Yokoyama	44		
<i>Neverita (Glassaulax) reiniana</i> (Dunker)	1	10-50	31-37
<i>Cryptonatica janthostomoides</i> (Kuroda and Habe)	10	20-300	31-43
<i>Siphonalia cf. fusoides</i> (Reeve)	1	10-100	32-41
<i>Fusinus perplexus</i> (A. Adams)	1	10-100	31-42
<i>Mitra</i> sp.	1		
<i>Fulgoraria cf. masudae</i> Hayasaka	1		
<i>Turbonilla</i> sp.	4		
Scaphopoda			
<i>Dentalium (Antalis) weinkauffi</i> (Dunker)	25	15-550	30-42

autochthonous or semiautochthonous. The number of specimens observed on the surface area of the block illustrated in Figure 7 is 13/222 cm², which is equivalent to a density of 585 ind./m². This value is nearly the same as that reported for Recent faunas (Fujita, 1992).

Together with ophiuroids, we found 23 molluscan species at this locality (Table 1). Species typical mainly of the lower sublittoral zone, such as *Limopsis (Empleconia) cumingi* and *Cryptonatica janthostomoides*, are common in the Hachioji Formation. The mode of occurrence of ophiuroids and molluscs suggests that the ophiuroids lived on a lower sublittoral, sandy silt bottom. Recent *O. s. sarsii* is distributed mainly on the uppermost continental slope around Japan (Fujita and Ohta, 1990). The Hachioji specimens lived in a slightly shallower setting than their extant counterparts. Some molluscs, e.g. *Anadara (Scapharca) ommaensis*, *Yoldia (Cnesterium) notabilis* and *Turritella (Neohaustator) saishuensis* have been recognized as typical of the Omma-Manganji fauna (Ogasawara, 1994), which suggests the sedimentary environment of the Hachioji Formation to have been one of a mild to cool temperate marine climate.

Acknowledgments

The authors would like to thank Masae Omori, Prof. Emeritus of Azabu University for identification of molluscan fossils and valuable comments. Our deep gratitude is

expressed to Iwao Kobayashi of Niigata University, Toshihiko Fujita of National Science Museum, Tokyo, John W. M. Jagt of Natuurhistorisch Museum, Maastricht, Netherlands, Frederick H.C. Hotchkiss of the Panometrics Company, U.S.A. and Gordon Hender of Natural History Museum of Los Angeles County for critical reading of the manuscript. Our special thanks are due to the following persons for their kind help in various ways: Tsunemi Kubodera of National Science Museum, Tokyo, Hideaki Nagamori of Niigata University, Akiko Nagao and Matthew Willemssen of Hitotsubashi Senior High School and Masaki Kajihara of Tate Senior High School. We wish to thank the Showa Seitoku Memorial Foundation for their generous financial assistance during 1997 and 1998.

References cited

- Arato, H., 1997: The Plio-Pleistocene stratigraphy of the Kanbara Plain, Niigata Prefecture, Central Japan and its sequence stratigraphic correlation with the Uonuma Group. *Earth Science*, vol. 51, no. 3, p. 233-244.
- Berry, C.T., 1934: Miocene and Recent *Ophiura* skeletons. *Johns Hopkins University Studies in Geology*, vol. 11, p. 9-135.
- Bjørlykke, K.O., 1898: Geologisk kart med beskrivelse over Kristiania by. *Norges Geologiske Undersøgelse*, no. 25, p. 1-86.

- Clark, H.L., 1911 : North Pacific ophiurans in the collection of the United States National Museum. *Bulletin of the United States National Museum, Smithsonian Institution*, vol. 75, p. 1-302.
- D'yakonov, A.M., 1954 : Ophiuroids of the USSR seas. *Zoological Institute of the Academy of Science of the USSR*, No. 55, p. 1-123. (in Russian, translated, 1967 : Israel Program for Scientific Translations)
- Fujita, T., 1992 : Dense beds of ophiuroids from the Paleozoic to the Recent : the significance of bathyal populations. *Otsuchi Marine Research Center Report*, No. 18, p. 25-41.
- Fujita, T. and Ohta, S., 1989 : Spatial structure within a dense bed of the brittle star *Ophiura sarsi* (Ophiuroidea : Echinodermata) in the bathyal zone off Otsuchi, north-eastern Japan. *Journal of the Oceanographical Society of Japan*, vol. 45, p. 289-300.
- Fujita, T. and Ohta, S., 1990 : Size structure of dense populations of the brittle star *Ophiura sarsii* (Ophiuroidea : Echinodermata) in the bathyal zone around Japan. *Marine Ecology Progress Series*, vol. 64, p. 113-122.
- Habe, T., 1977 : *Systematics of Mollusca in Japan, Bivalvia and Scaphopoda*, 372 p. Hokuryukan, Tokyo. (in Japanese)
- Higo, T., 1973 : *A general catalogue of the marine molluscan faunas around the Japanese Islands*, 397 p. Biological Society of Nagasaki. (in Japanese)
- Irimura, S., 1990 : Ophiuroidea. *Echinoderms from Continental Shelf and Slope around Japan*, Volume 1, p. 65-100. Tokyo, Japan Fisheries Resource Conservation Association.
- Ishida, Y. and Fujita, T., 1998 : *Ophiura sarsii sarsii* Lütken from the Middle-Late Miocene Wakkanai Formation in Hokkaido. *Palaeontological Society of Japan, 1998 Annual Meeting, Abstract*, p. 93. (in Japanese)
- Ishida, Y. and Inoue, K., 1993 : Fossil ophiuroids (*Ophiura sarsii*, Lütken, 1854) from the Pleistocene Ichijuku Formation (Kazusa Group), Chiba Prefecture, Central Japan. *Natural History Research*, vol. 2, No. 2, p. 103-112.
- Ishida, Y. and Inoue, K., 1995 : Mode of occurrence and paleoecology of *Ophiura sarsii* Lütken, 1854-Fossil ophiuroids from the Pleistocene Ichijuku Formation (Kazusa Group), Chiba Prefecture, Central Japan-. *Earth Science*, vol. 49, p. 32-44.
- Ishida, Y., Kurita, Y., Nagamori, H. and Narita, T., 1997 : Fossil *Ophiura sarsii sarsii* from the Pliocene Hachioji Formation in Kashiwazaki City, Niigata Prefecture and the Upper Miocene Ogawa Formation from Shinshu-shinmachi, Nagano Prefecture. *Palaeontological Society of Japan, 1997 Annual Meeting, Abstract*, p. 19. (in Japanese)
- Ishida, Y., Nagamori, H. and Narita, K., 1998 : *Ophiura sarsii sarsii* (Echinodermata, Ophiuroidea) from the Late Miocene Ogawa Formation, Shinshushinmachi, Nagano Prefecture, Central Japan. *Research Report of the Shinshushinmachi Fossil Museum*, no. 1, p. 9-16.
- Ishida, Y., Tanabe, T., Ito, T. and Hachiya, K., 1996 : Fossil ophiuroids from the Plio-Pleistocene Hijikata Formation of the Kakegawa Group, Shizuoka, Central Japan. *Bulletin of the National Science Museum, Tokyo, Series C*, vol. 22, nos. 3, 4, p. 63-89.
- Ishida, Y., Tokairin, H. and Nagasawa, K., 1997 : Fossil ophiuroids from the Upper Miocene Hongo Formation in Asahi-machi, Yamagata Prefecture, Northern Japan. *Geological Society of Japan, 104th Annual Meeting, Abstract*, p. 355. (in Japanese)
- Jensen, M. and Thomsen, E., 1987 : Ultrastructure, dissolution and "pyritization" of Late Quaternary and Recent echinoderms. *Bulletin of the Geological Society of Denmark*, vol. 36, p. 275-287.
- Kobayashi, I., Tateishi, M., Kurokawa, K., Yoshimura, T. and Kato, H., 1989 : *Geology of the Okanomachi District*. 112 p. With Geological Sheet Map at 1 : 50,000, Geological Survey of Japan. (in Japanese with English abstract)
- Kuroda, T. and Habe, T., 1952 : *Checklist and bibliography of the Recent marine Mollusca of Japan*. 210 p. Zoological Institute, College of Science, Kyoto University.
- Lütken, C.F., 1855 : Bidrag til Kundskab om Slangestjernerne. 1. Forelobig Oversigt over Gronlandshavets Ophiurer. *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i Kjobenhavn*, 1854, p. 95-104.
- Matsumoto, H., 1917 : A monograph of Japanese Ophiuroidea, arranged according to a new classification. *Journal of the College of Science, Imperial University Tokyo*, vol. 38, article 2, p. 1-408, 52 pls.
- Mortensen, T., 1927 : *Handbook of the Echinoderms of the British Isles*, 471 p., 269 figs. London, Oxford Univ. Press.
- Ogasawara, K., 1994 : Neogene paleogeography and marine climate of the Japanese Islands based on shallow-marine molluscs. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 108, p. 335-351.
- Yasui, S., Kobayashi, I. and Tateishi, M., 1983 : Stratigraphy of the Uonuma Formation in the Hachikoku Oil Field and the Southern part of Chuo Oil Field, Niigata Prefecture, Central Japan. *Earth Science*, vol. 37, no. 1, p. 22-37. (in Japanese with English abstract)